Impact of climate change on the French part of the River Meuse -the CHIMERE 21 project

Meuse symposium 21 September 2021

Context and goals of the project

The CHIMERE 21 project

CHIers - MEuse: hydrological Regime Evolution in the 21st century

Topic of the project: Study of the impact of climate change on the streamflows of the Chiers and Meuse Rivers

Partners: INRAE (formerly Irstea), Météo-France, EDF, Université de Lorraine, DREAL Grand-Est

Funding by Agence de l'Eau Rhin-Meuse



The context

CHIers - MEuse: hydrological Regime Evolution in the 21st century

Jacob et al. (2013)

Global scale evolutions need to be refined at the local scale through specific studies

Past studies:

- Explore 2070: France-wide project
- Amice: Meuse-wide project

Intermediate scenario(P)



Intermediate scenario (T)



Scenario without mitigation (P)



Scenario without mitigation (T)



Mean evolution of precipitation (left) and air temperature (right) by 2071-2100

Objectives of CHIMERE 21

- Update existing knowledge (Explore 2070 and Amice are beginning to be old)
- Refine results (needs of local studies)
- Huge stakes for low flows

Objectives of the project:

- Study of the impact of climate change on future Meuse streamflows
- Focus on uncertainties
- Production of synthetic sheets

Presentation of the basin and data

Hydro data





16 stations (one for around 480 km²)

- Dubious daily discharge values removed through visual inspection
- At Chooz, the evaporation water consumption was estimated using daily air temperature and the nuclear power plant charge
- Other influences were not removed from time series:
  - Either low impact at the basin scale
  - Or too much uncertainty so data were just discarded

Observed meteo data

Météo-France SAFRAN reanalysis: combination between observed in situ data and model simulations

- Spatialised data on a regular grid: 8 km x 8 km
- Daily data

- Potential evapotranspiration = Penman-Monteith (using SAFRAN variables)



Climate change over the Meuse basin

Climate change over the Meuse basin

Modelling chain: composed of several steps

RCPs



Source : Knutti and Sedlácek (2013)

GCMs



Source : Climate Change 2014: Impacts, Adaptation, and Vulnerability. Part A: Global and Sectoral Aspects

RCMs



Source : Jacob et al., 2013

Corrections de biais/
Descente d'échelle



Source : Drias. Exemple méthode quantile-quantile

Modélisation hydrologique



Echelle d'étude : le bassin versant

Cascade d'incertitude



Wilby and Dessai (2010)

Figure from Lemaitre-Basset (2020)

Climate change over the Meuse basin

Selection of climate data for CHIMERE 21:

RCPs



Source : Knutti and Sedlácek (2013)

GCMs



Source : Climate Change 2014: Impacts, Adaptation, and Vulnerability. Part A: Global and Sectoral Aspects

RCMs



Source : Jacob et al., 2013

.RCP 4.5

.RCP 8.5

	GCM	RCM
Couple 1	CNRM-CM5	ALADIN53
Couple 2	IPSL-CM5A	IPSL-INERIS-WRF
Couple 3	CNRM-CM5	RCA4
Couple 4	IPSL-CM5A	RCA4
Couple 5	MPI-ESM	RCA4

3 different GCMs, 3 different RCMs

Climate change over the Meuse basin

Seasonal climate evolution

Large uncertainty

Summer (all models and RCPs)	Temperatures	Precipitation
Near future	+0,4 to +1,2 °C	-14 to +23 %
Far future	+1 to +4,4 °C	-39 to +21 %

Winter (all models and RCPs)	Temperatures	Precipitation
Near future	+0,8 to +1,5 °C	+1 to +35 %
Far future	+1,8 to +4,4 °C	+18 to +57 %

« Dry » summer: 2 couples

« Neutral » summer: 1 couple

« Wet » summer: 2 couples

Far future
RCP 8.5

Difference related to the historical period







INRAE METEO FRANCE EDF UNIVERSITÉ DE LORRAINE DIRECTION RÉGIONALE DE L'ENVIRONNEMENT, DE L'AMÉNAGEMENT ET DU LOGEMENT GRAND EST

The CHIN
21 Septe

Hydrological modelling

Hydrological modelling

RCPs

Source : Knutti and Sedlácek (2013)

GCMs

Source : Climate Change 2014: Impacts, Adaptation, and Vulnerability. Part A: Global and Sectoral Aspects

RCMs

Source : Jacob et al., 2013

Corrections de biais/
Descente d'échelle

Source : Drias. Exemple méthode quantile-quantile

Modélisation hydrologique



Echelle d'étude : le bassin versant

Cascade d'incertitude

Wilby and Dessai (2010)

Figure from Lemaitre-Basset (2020)

Hydrological modelling

Four hydrological models

Several models are necessary to verify to which extent all models provide similar trajectories

GRSD

Conceptual
Semi-distributed
(sub-basins)

MORDOR

Conceptual
Semi-distributed
(sub-basins)

PRESAGES

Conceptual
Lumped

SIM2

Physically-based
Distributed
(regular grid)

1976 Drought



2003 drought



Impact of climate change on hydrology

Impact of climate change on hydrology

RCPs

Source : Knutti and Sedlácek (2013)

GCMs

Source : Climate Change 2014: Impacts, Adaptation, and Vulnerability. Part A: Global and Sectoral Aspects

RCMs

Source : Jacob et al., 2013

Corrections de biais/
Descente d'échelle

Source : Drias. Exemple méthode quantile-quantile

Modélisation hydrologique

Echelle d'étude : le bassin versant

Cascade d'incertitude



Wilby and Dessai (2010)

Future regimes

RCP 8.5, Horizon 2071-2100

Analysis of future regimes (all climate models included) at Chooz

Large increase of streamflows during the high flow period

Large uncertainty of low flows



Evolution of mean flows along the Meuse

RCP 4.5



RCP 8.5





RCP 4.5: increase of mean flows, especially downstream

RCP 8.5: increase everywhere, especially for the far future

Evolution of high flows along the Meuse

RCP 4.5



RCP 8.5





RCP 4.5: moderate increase (only for the far future)
RCP 8.5: important increase, especially downstream

Evolution of low flows along the Meuse

RCP 4.5



RCP 8.5





RCP 4.5: slight decrease, especially upstream

RCP 8.5: moderate decrease upstream for far future, slight otherwise

Hierarchy of uncertainties for low flows

Indicator = VCN3

Internal variability dominates for near future

GCMs and RCMs dominate then

Other sources remain limited



Probabilities of different trends

Need to answer the following questions: « What is the probability that an indicator change of X % under climate change? » et « At which temporal horizon could this trend emerge? »



Pourcentage of all simulations indicating a trend of at least 10% of low flows(VCN3)

Conclusions and perspectives

Main conclusions of CHIMERE 21

Hydroclimate evolutions

Climate:

- Temperature increase, especially for RCP 8.5 and far future
- Heterogeneous precipitation evolution: increase for winter, large uncertainty for summer
- Projections of CHIMERE 21 warmer and wetter than Explore 2070 but consistent with Drias 2020

Hydrology:

- Large uncertainties for summer, possible decrease of streamflows for upstream
- Increase during winter, especially downstream
- Climate brings most of uncertainties on Meuse streamflows

Perspectives

The CHIMERE 21 study gave thoughts for food regarding adaptation strategies

- Evolutions of streamflows question strong stakes:
  - Increase of streamflow downstream, which is already identified as a territory with high risk of floods
  - Possible decrease of low flows upstream

Explore 2: a France-wide project aiming at evaluating impacts of climate change using Drias 2020 (started recently)

LIFE Eau&Climat (https://www.gesteau.fr/life-eau-climat): a project aiming at helping local water stakeholders to evaluate climate change impacts, to take them into account for planning and to undertake adaptation strategies

Bonus

Synthetic sheets

Projet CHIMERE 21

Fiche synthétique

B7200000 – La Meuse @ Chooz [Trou du Diable]



<- page 1

Page 2 ->

CALAGE DES MODELES

PERFORMANCE SUR 1970-2016

Calage régime





Calage étiages





ROBUSTESSE



PROJECTIONS FUTURES

Régimes interannuels projetés, tous modèles hydrologiques confondus : période de référence = 1976-2005, périodes futures = 2021-2050 (horizon 2035) et 2071-2100 (horizon 2085)



Incertitudes et tendances



LEGENDES



Pour plus de détails sur la méthodologie et l'interprétation des résultats,
Thirel, G., Collet, L., Rousset, F. et al. (2021).
Rapport final du projet CHIMERE 21, 152 p. https ://hal.archives-ouvertes.fr/hal-03206168.

Contact : guillaume.thirel@inrae.fr
Produit par : Lila Collet et Guillaume Thirel

Le 23 avril 2021

The CHIMERE 21 team

Guillaume Thirel
(INRAE Antony, coordination)

Lila Collet
(INRAE Antony)

Fabienne Rousset
(Météo-France)

Olivier Delaigue
(INRAE Antony)

Joël Gailhard
(EDF)

Matthieu Le Lay
(EDF)

Didier François
(Université de Lorraine)

Charles Perrin
(INRAE Antony)

Mathieu Reverdy
(INRAE Lyon)



Raphaëlle Samacoits
(Météo-France)

Morgane Terrier
(INRAE Antony)

Jean-Philippe Vidal
(INRAE Lyon)



Jean-Pierre Wagner
(DREAL Grand-Est)