



[www.openda.org](http://www.openda.org)


# Calibration of an estuary with Delft-3D flow

**Martin Verlaan, Firmijn Zijl**

**Stef Hummel, Julius Sumihar, Nils van Velzen,  
Albrecht Weerts, Ghada El Serafy, Herman  
Gerritsen, ...**



**Deltares, TU Delft, VORtech**

# Installation Delft3d v4 and OpenDA v2.1 for Windows




- Unpack delft3d license file (and remember where to)
- Start delft3d installer and install all items ( skip manuals )
  - Give default answers everywhere
  - Select license file that you just unpacked when asked
- Make a shortcut for quickplot to the desktop (from C:\Program Files (x86)\Deltares\Delft3D 4.01.00.rc.09\win32\quickplot\bin)
- Unpack openda v2.1 for windows to [c:\openda](#) (or somewhere else as long as it has no spaces in the path)
- Unpack the delft3d-plugin for openda to the same location.  
NOTE: the bin-folders and their contents should merge to the same place.
- Make a shortcut for oda\_rungui.bat to de desktop

# Simplified model of an estuary




River discharge boundary

- One dimensional model
- Tidal boundary M2 (12h25min) and S2 (12h)
- Constant slope depth
- Constant river inflow
- 3 Observation locations
- Observations are not real but generated with 'truth' model.





# Questions




- Download estuary.zip from <http://www.openda.org/course> and unpack
- Install openda\_d3d\_plugin.zip in the OpenDA bin-directory
- Run the simulation with OpenDA, using the main OpenDA file simulate.oda
- Prepare some time-series plots with quickplot
  - Start matlab in direcory src/tools\_lgpl/matlab/quickplot/progrsrc and run d3d\_qp
  - the observations are available as tekal file, for including them in the plots (use add to plot and change the color)
  - Output can be found in estuary/work/work0
- What are the most likely causes of differences between observations and model?

# Initial performance




RMSE=9cm

# Initial performance





water level (m)




RMSE=12cm

# Initial performance



RMSE=7cm


# Questions




- Run the calibration for a globally constant change to the bathymetry (experiment DEP)
  - Start OpenDA with estuary/calibration.oda
  - Look at the output in the control tab and output tab
  - The output of each of the runs can be found in work/work<number>
  - Plot the time-series with quickplot.
- Is this what you expected?



# Calibration Depth





## Station 1





Name	First guess	DEP calib.
Station 1	9.2 cm	4.5
Station 2	12.7	5.0
Station 3	7.1	4.9
Cost	574	134


# DEP output Station 2



# DEP output Station 3




# Questions




**Since the amplitude shows a similar deviation from observations in the whole domain we add calibration of M2 tides at boundary.**

- Add the calibration (experiment DEP+M2)
  - Uncomment M2 section in stochModel/D3DStochModel.xml
  - Run calibration
  - Look at the output and plot the time-series.
  - Is this what you expected?

# Calibration Depth+M2





Station 1





Name	First guess	DEP	DEP+M2
Station 1	9.2 cm	4.5	1.1
Station 2	12.7	5.0	1.4
Station 3	7.1	4.9	1.1
Cost	574	134	8.4


# DEP+M2 output Station 2



# DEP+M2 output Station 3



# Questions





**The output looks nice. The cost-function is much lower, but there is still a problem...**

- Make a longer run with the final run of experiment DEP+M2
  - Modify `work/work<last_number>/estuary.mdf` and change the `Tstop = 2.3040000e+004` which is 17-1-1991 0:00h; alternatively use the `delft3d-gui`.
  - Run `deltares_hydro.exe` for this case
  - Make time-series plots
  - What is wrong?



# Long run for DEP+M2 result




# Questions

**The error in S2 was attributed to M2. Let's make fix this with a longer simulation and adding S2 to the calibration**

- Add S2 to calibration and lengthen simulation experiment DEP+M2
  - Modify input\_d3d/estuary.mdf and change the Tstop = 2.3040000e+004 which is 17-1-1991 0:00h; see also estuary\_long.mdf
  - Lengthen the observations in stochobserver/noosObservations.xml to 17-1-1009 0:00h; see noosObservations\_long.xml
  - Uncomment S2 section in stochModel/D3DStochModel.xml
  - Run calibration with OpenDA
- What would go wrong if we would use only 3 days of observations for calibration of S2 and M2?


# Calibration DEP+M2+S2



Name	First guess	DEP+M2+S2
Station 1		0.9cm
Station 2		0.7
Station 3		0.2
Cost	5281	1.5

Parameter	Final value (change)	True values
M2.Amplitude	0.1 cm	0.0 cm
M2.Phase	0.4 degr	0.0 degr
S2.Amplitude	10.1 cm	10.0 cm
S2.Phase	0.3 degr	0.0 degr
Depth	-92cm	-100cm

# And much more



- Calibration of roughness
- Calibrate blocks of the grid for depth or roughness
- Proportional instead of additive modification of parameters
- Make subselections of observations
- Restarts
- Parallel computing
- Output formats and selection
- Try other algorithms
- Calibration of other models, such as sobek, swan or waqua
- ....